

## 10: The Derivative and the Tangent Line

### Key Terms

- **Secant Line:** A secant line is a line that passes through two points that lie on a curve.
- **Tangent Line:** A tangent line to a curve at a given point is a line that barely touches the curve at that point.
- **Average Rate of Change:** The average rate of change is the average rate at which a function changes over a given interval. It is represented by the slope of a secant line.
- **Instantaneous Rate of Change:** The instantaneous rate of change is the rate of change of the function at a single point. It is represented by the slope a tangent line.
- **Derivative:** The derivative of a function  $f(x)$  is another function  $f'(x)$  which gives the slope of the tangent line for any value of  $x$ .


### Definition of the Derivative

The derivative, denoted by  $f'(x)$  gives the slope of the tangent line to a curve  $f(x)$ . Since the secant line approaches the tangent line when  $h \rightarrow 0$ , the slope of the secant line (i.e., the difference quotient) approaches the slope of the tangent line (i.e., the derivative) when  $h \rightarrow 0$ .

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$


### Interpretation of the Derivative

Suppose that  $f(x) = x^2$ . The derivative is  $f'(x) = 2x$ . Now consider the point  $(-2, 4)$  on the graph of  $f(x)$ . The slope of the tangent line at the point  $(-2, 4)$  would be given by  $f'(-2) = 2(-2) = -4$ .



### Tangent Line


The secant lines converge to the tangent line as the two points get closer and closer.



### The Difference Quotient

The difference quotient is defined to be the slope of the secant line:

$$\text{Slope} = \frac{f(x_0 + h) - f(x_0)}{h}$$



### Example

Example: Find the derivative of  $f(x) = -2x^2$ .

Solution:  
The derivative is:

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-2(x+h)^2 - (-2x^2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-2(x^2 + 2xh + h^2) + 2x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{-2x^2 - 4xh - 2h^2 + 2x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{\cancel{-2x^2} - 4xh - 2h^2 \cancel{+ 2x^2}}{h} \\ &= \lim_{h \rightarrow 0} \frac{-4xh - 2h^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(-4x - 2h)}{h} \\ &= \lim_{h \rightarrow 0} (-4x - 2h) \\ &= -4x \end{aligned}$$

### Horizontal Tangent Line

A tangent line is horizontal whenever the derivative equals zero.

### Continuity

If the derivative of a function exists at a point, the function is guaranteed to be continuous at that point.

How to Use This Cheat Sheet: These are the keys related this topic. Try to read through it carefully twice then rewrite it out on a blank sheet of paper. Review it again before the exams.